

AD-A049 693

PRINCETON UNIV N J DEPT OF STATISTICS
LIMIT THEOREMS FOR THE SIMPLE BRANCHING PROCESS ALLOWING IMMIGR--ETC(U)
DEC 77 A G PAKES, A D BARBOUR
TR-134-SER-2

F/G 12/1

N00014-75-C-0453

NL

UNCLASSIFIED

| OF |

AD
A049 693



END
DATE
FILMED
3-78
DDC

AD A 049693

AD No. JDC FILE COPY

2



**LIMIT THEOREMS FOR THE SIMPLE BRANCHING
PROCESS ALLOWING IMMIGRATION, II: THE
CASE OF INFINITE OFFSPRING MEAN**

by

**A. D. Barbour
Cambridge University**

and

**Anthony G. Pakes*
Princeton University**



**Technical Report No. 134, Series 2
Department of Statistics
Princeton University
December 1977**

See 1479
in
back

Research sponsored in part by a contract with the
Office of Naval Research, No. N00014-75-C-0453, awarded
to the Department of Statistics, Princeton University.

*On leave from the Dept. of Mathematics,
Monash University, Clayton, Victoria,
Australia.

A B S T R A C T

This paper obtains some limit theorems for the simple branching process allowing immigration $\{X_n\}$ when the offspring mean is infinite. It is shown that there exists a function U such that $\{e^{-n}U(X_n)\}$ converges almost surely and if $s = \sum b_j \log^+ U(j) < \infty$, where $\{b_j\}$ is the immigration distribution, the limit is non-defective and non-degenerate but is infinite if $s = \infty$.

When $s = \infty$ limit theorems are found for $\{U(X_n)\}$ which involve a slowly varying non-linear norming.

Key words: Bienaymé-Galton-Watson branching process, immigration, martingale convergence, limit theorems, regular variation.

ACCESSION for	
NTIS	Write Section <input checked="" type="checkbox"/>
DDC	B&H Section <input type="checkbox"/>
UNANNOUNCED	
JUSTIFICATION	
BY	
DISTRIBUTION/AVAILABILITY CODES	
Dist.	AIL and/or SPECIAL
A	

1. INTRODUCTION

Let $\{X_n: n=0,1,\dots\}$ denote a Bienaymé-Galton-Watson process with immigration, for which the offspring probability generating function, f , satisfies $m = f'(1-) = \infty$ and $\{I_n: n=1,2,\dots\}$, the number of immigrants into successive generations, are independent and identically distributed with probability generating function b satisfying $b(0) < 1$. Suppose also that $X_0 = 0$.

This paper describes the behavior of X_n as $n \rightarrow \infty$ in terms of appropriate limit theorems. In §2 we show that for a certain increasing function U constructed as in [6], the sequence $\{e^{-n}U(X_n)\}$ converges almost surely and that the limit is non-defective if a certain condition on the immigration distribution obtains and is essentially infinite otherwise. It is pointed out that, in the former case, $\{X_n\}$ may be classified in terms of regularity and irregularity exactly as in the case of no immigration [6].

In §3 we consider the case where $Y_n = e^{-n}U(X_n) \rightarrow \infty$ a.s. Limit theorems are obtained for Y_n which involve a non-linear norming by a slowly varying (SV) function. Theorem 2 involves no extra assumptions but has a rather strange appearance: neater versions are given in Corollaries 1-3 under conditions expressed in terms of $G(x) = 1 - b(\exp[-1/V(e^x)])$, where V is the functional inverse of U . These results are analogues of the families of limit theorems for the case $m < \infty$ which were presented in Part I [3], and some examples are given to show that the conditions in Corollaries 1-3 are not vacuous. Finally, it is shown that there is no sequence (c_n) of norming constants such that Y_n/c_n has a limit in distribution which is neither defective nor degenerate at the origin.

To complete the survey in Part I, it suffices to mention that a version of Theorem 1 was first proved in [2] under conditions on f ensuring that U could be chosen as a suitable power of \log^+ and only convergence in law was established. The conditions of this result were substantially relaxed in [1] to allow U to be SV at infinity and strictly increasing. A restricted and more opaque version of Corollary 1 was proved in [2].

2. ALMOST SURE CONVERGENCE

Let f_n denote the n^{th} functional iterate of f and h_n the functional inverse of $-\log f_n(e^{-s})$. Let $p_n(s) = \prod_{m=1}^n b(\exp(-h_m(s)))$ and $p(s) = \lim_{n \rightarrow \infty} p_n(s)$ for $0 < s < -\log q$ where $q = f(q) < 1$. Since $h_n(s) \downarrow 0$ ($n \rightarrow \infty$) it is clear that either $p(s) > 0$ or $\equiv 0$. In the former case it is clear that $\{[\exp(-h_n(s)X_n)]/p_n(s)\}$ is a martingale and the martingale convergence theorem shows that for each $0 < s < -\log q$, $\exp(-h_n(s)X_n) \xrightarrow{\text{a.s.}} W(s)$, say, and $E W(s) = p(s)$. The behavior of $\{X_n\}$ can be analyzed in precisely the same way in which that of the corresponding process without immigration was analyzed in [6]. In particular the classification of points in $(0, -\log q)$ as regular and irregular carries over to the present case and if $T = \sup\{s | 0 < s < -\log q, W(s) < 1\}$ it follows that T is non-defective and $P(T \geq s_r) = p(s_r)$, $P(T = s_r) = 0$ for every regular point s_r .

Finally, for the function U constructed in [6], the arguments used in this reference show that $e^{-nU(X_n)} \xrightarrow{\text{a.s.}} U(T^{-1})$. In the complementary case, when $p(s) \equiv 0$, the martingale convergence theorem shows that $\exp(-h_n(s)X_n) \xrightarrow{\text{a.s.}} 0$ for each $0 < s < -\log q$. To proceed

further we need the following details about U . It is shown in [6] that by starting with a fixed $s_0 \in (0, -\log q)$ it is possible to construct U so that it is continuous on $[0, \infty)$, identically zero on $[0, 1/(-\log q)]$, positive and strictly increasing to infinity on $(1/(-\log q), \infty)$ and satisfies the relations

$$U(1/h_n(s_0)) = e^n, \quad U(1/h_n(s))/U(1/h_n(s_0)) = U(1/s) \quad (1) \\ (n=0, 1, \dots)$$

Choose a sequence $\{s_m; m=1, 2, \dots\}$ such that $-\log q > s_m + 0 \quad (m \rightarrow \infty)$. Let Ω_m be a subset of the basic probability space such that $P(\Omega_m) = 1$ and $h_n(s_m)X_n(\omega) \rightarrow \infty \quad (n \rightarrow \infty; \omega \in \Omega_m)$. Thus on $\Omega' = \bigcap_{m \geq 1} \Omega_m$, $h_n(s_m)X_n(\omega) \rightarrow \infty$ for each m and this implies that eventually $X_n(\omega) \geq 1/h_n(s_m)$ and hence, from (1), eventually

$$e^{-n}U(X_n(\omega)) \geq U(1/s_m) \quad (\omega \in \Omega').$$

Letting $n \rightarrow \infty$ and then $m \rightarrow \infty$, we obtain all except the final assertion of the following theorem.

Theorem 1. If $p(s) > 0$ then $e^{-n}U(X_n) \xrightarrow{a.s.} U(1/T)$ where T is defined above, and if $p(s) = 0$, $e^{-n}U(X_n) \xrightarrow{a.s.} \infty$. Furthermore, $p(s) > 0$ iff

$$\sum_{j=1}^{\infty} b_j \log^+ U(j) < \infty \quad (2)$$

where $\{b_j\}$ denotes the immigration distribution.

The last assertion follows from these observations. Since U is strictly increasing, the relations (1) can be solved to obtain an explicit expression

$$h_n(s) = 1/V(e^n U(s^{-1})) \quad (3)$$

for h_n , and the proof of Lemma 2 in [1] needs, with one exception,

only trivial changes to show that (2) holds iff

$$\prod_{n=1}^{\infty} b\left(\exp(-1/V(e^n U(s^{-1})))\right) > 0$$

where V is the functional inverse of U , whence the assertion in this case. The exception is that the proof given in [1] assumes that $\log U$ is SV: that this is so follows from

Lemma 1. If L is SV at infinity then so is $L(U(\cdot))$.

Proof. By virtue of the uniform convergence theorem for SV functions [7] it suffices to show that for each $\lambda > 0$, $U(\lambda x)/U(x)$ is bounded away from zero and infinity for all sufficiently large x . It suffices to consider the case $1 \leq \lambda < \infty$. The proof to be given parallels that of Lemma 2.3.3 in [6]. Choose $s_0 \in (0, -\log q)$ as in Construction 2.3.1 of [6], and set $D = [h(s_0), s_0]$. The functions $g_n(s) = h_{n+1}(s)/h_n(s)$ are continuous on D , and, from equations (2) and (3) of [6], it follows that

$$g_n(s) \geq g_{n+1}(s) ; \quad g_n(s) \rightarrow 0 \quad \text{as } n \rightarrow \infty : \quad (4)$$

hence, by Dini's theorem, this convergence is uniform on D . Thus for all $n \geq n_0$, chosen large enough, and all $s \in D$, $\lambda/h_n(s) \leq 1/h_{n+1}(s)$. For each sufficiently large x construct n and $s \in D$ so that $x^{-1} = h_n(s)$ (see Construction 2.3.1, [6]). It follows that

$$1 < U(\lambda x)/U(x) \leq U(1/h_{n+1}(s))/U(1/h_n(s)) = e ,$$

and the lemma now follows.

3. THE CASE $p(s) \equiv 0$

The description of the behavior of X_n in the case $p(s) \equiv 0$ is through distributional, rather than almost sure, limit theorems.

The first step in deriving them is to note that, from (3),

$$E\left(\exp(-h_n(t)X_n)\right) = p_n(t) = \prod_{m=1}^n b\left[\exp\left(-1/V(e^m U(t^{-1}))\right)\right].$$

Using integral test comparisons, as in [3], it is not difficult to show that

$$p_n(t) = \Delta_n(t) \exp \int_0^n \log b\left[\exp\left(-1/V(e^y U(t^{-1}))\right)\right] dy$$

where $\Delta_n(t_n) \rightarrow 1$ if $t_n \rightarrow 0+$ as $n \rightarrow \infty$. Denoting the integral by J_n and making the change of variable $e^x = e^y U(t^{-1})$, it follows that if $t = t_n \rightarrow 0+$ then

$$-J_n \sim J_n = \int_{\log U(1/t_n)}^{n + \log U(1/t_n)} G(x) dx \quad (5)$$

where $G(x) = 1 - b\left[\exp(-1/V(e^x))\right]$. We are now in a position to approach the main limit theorem, by making a suitable choice of t_n , and by using the following version of Reuter's Lemma 1 in [1].

Lemma 2. Let U be constructed according to Construction 2.3.1 in [6] and in addition suppose it is strictly increasing on $(1/(-\log q), \infty)$.
Let y_n be positive, increasing and satisfy the properties

(a) $y_n(u)/y_n(v) \rightarrow \infty$ ($n \rightarrow \infty$) if $u > v$ and $0 < c < u, v < d \leq \infty$; and

(b) $e^n y_n(u) \rightarrow \infty$ ($n \rightarrow \infty$; $c < u < d$).

If, for a sequence of non-negative random variables $\{W_n\}$ there is a continuous function $a(\cdot)$ such that

$$E\left[\exp\left(-W_n/V(e^n y_n(u))\right)\right] \rightarrow a(u) \quad (c < u < d; n \rightarrow \infty)$$

then

$$P(e^{-n} U(W_n) \leq y_n(u)) \rightarrow a(u).$$

Proof. Let $A_n = \{e^{-n}U(W_n) \leq y_n(u)\} = \{W_n \leq V(e^n y_n(u))\}$. Choose $u \in (c, d)$ and $c < u_1 < u < u_2 < d$. If $Y_n^{(i)} = \exp\{-W_n/V(e^n y_n(u_i))\}$ ($i=1,2$), then arguing as in [1] we obtain

$$EY_n^{(1)} - \exp(-\lambda_n^{(1)}) \leq P(A_n) \leq (EY_n^{(2)}) \exp \lambda_n^{(2)}$$

where $\lambda_n^{(i)} = V(e^n y_n(u))/V(e^n y_n(u_i))$ ($i=1,2$). If we can show that $\lambda_n^{(1)} \rightarrow \infty$ and $\lambda_n^{(2)} \rightarrow 0$, the assertion follows upon then letting $u_i \rightarrow u$.

We shall prove only that $\lambda_n^{(1)} \rightarrow \infty$. It follows from (a) that for each $t > 1$

$$\lambda_n^{(1)} \geq V(te^n y_n(u_1))/V(e^n y_n(u_1))$$

for all sufficiently large n . Let $m(n)$ be the integer part of $n + \log y_n(u_1)$. By (b) $m(n) \rightarrow \infty$ ($n \rightarrow \infty$) and $e^{m(n)} \leq e^n y_n(u_1) \leq e^{m(n)+1}$ and hence, taking $t = e^2$,

$$\lambda_n^{(1)} \geq V(\exp\{m(n)+2\})/V(\exp\{m(n)+1\}) = h_{m(n)}(1/V(e))/h_{m(n)+1}(1/V(e)),$$

where we have used (3). Hence, from (4), $\lim_{n \rightarrow \infty} \lambda_n^{(1)} = \infty$.

Now, for $x \geq 1$, let $\Lambda(x) = \exp \int_0^{\log x} G(y) dy$, which is SV at infinity and strictly increasing. Furthermore, the ratio

$$\begin{aligned} \Lambda(x)/\Lambda(xe^n) &= \exp \left\{ - \int_{\log x}^{\log x+n} G(y) dy \right\} \\ &= \exp \left(- \int_1^{e^n} G(\log(zx)) dz/z \right) \end{aligned}$$

increases with x on $(1, \infty)$ from m_n say, to unity, and since $p(t) \equiv 0$ iff $\int_0^\infty G(x) dx = \infty$, it follows that $m_n \rightarrow 0$ ($n \rightarrow \infty$); moreover, for each fixed $x \geq 1$, $\Lambda(x)/\Lambda(xe^n)$ decreases to zero as $n \rightarrow \infty$. Hence, for any $0 < u < 1$ and all n sufficiently large, we may define $y_n(u) > 0$ uniquely in such a way that

$$\Lambda(y_n(u)) = u\Lambda(e^n y_n(u)),$$

where it follows that y_n is increasing on $(0,1)$, that $y_n(u) \rightarrow \infty$ ($n \rightarrow \infty$) and that if $0 < v < u < 1$,

$$\frac{y_n(u)}{y_n(v)} = \frac{\Lambda^{-1}\left\{u\Lambda(e^n y_n(u))\right\}}{\Lambda^{-1}\left\{v\Lambda(e^n y_n(v))\right\}} \geq \frac{\Lambda^{-1}\left\{u\Lambda(e^n y_n(v))\right\}}{\Lambda^{-1}\left\{v\Lambda(e^n y_n(v))\right\}} \rightarrow \infty$$

as $n \rightarrow \infty$, since Λ is SV at infinity. Choosing $U(t_n^{-1}) = y_n(u)$ in (5) we see that $j_n = u$ and hence $E\left[\exp\left\{-X_n/v(e^n y_n(u))\right\}\right] \rightarrow u$ ($0 < u < 1$), and that the other conditions of Lemma 2 are satisfied. This completes the proof of the following theorem.

Theorem 2. If $p(s) \equiv 0$, then

$$\Lambda(e^{-n}U(X_n))/\Lambda(U(X_n)) \xrightarrow{d} W$$

where W is uniformly distributed on $[0,1]$.

Analogues of Theorem 2 also obtain when $m < 1$ and $1 < m < \infty$; see [3].

The proof of Theorem 2 has been carried out under the condition $X_0 = 0$. However

$$\begin{aligned} p_i^{(n)}(t) &= E\left[\exp(-X_n h_n(t)) | X_0 = i\right] \\ &= \left[f_n(\exp(-h_n(t)))\right]^i p_0^{(n)}(t) \\ &= e^{-it} p_0^{(n)}(t), \end{aligned}$$

and since t was chosen to converge to zero, we see that Theorem 2 is valid for any initial state. Furthermore, $\Lambda(e^{-n}U(\cdot))/\Lambda(U(\cdot))$ is strictly increasing and continuous and hence the weakly convergent sequence in Theorem 2 is a Markov chain and, in addition, is mixing [5, Theorem 2]. Thus [4] we cannot have convergence in probability

in Theorem 2.

The quantity converging in law in Theorem 2 has a rather odd appearance since $U(X_n)$ is present in both numerator and denominator. Tidier versions can be obtained by making estimates of

$$\Lambda(e^{-n}U(X_n))/\Lambda(U(X_n)) = \exp \left\{ -Y_n \int_{1-n/Y_n}^1 G(zY_n) dz \right\}; \quad (6)$$

under appropriate assumptions on the behavior of G . For brevity, we write $Y_n \equiv \log U(X_n)$ throughout.

Corollary 1. If $p(s) \equiv 0$ and $\lim_{x \rightarrow \infty} xG(x) = 0$, then

$$\Lambda(e^{-n}U(X_n))/\Lambda(e^n) \xrightarrow{d} W,$$

where W is uniformly distributed on $[0,1]$.

Proof. Since $xG(x) \rightarrow 0$, we have

$$\int_{Y_n - n}^{Y_n} G(y) dy = o \left\{ -\log(1-n/Y_n) \right\}; \quad (7)$$

combining Theorem 2, (6) and (7), it follows that, as $n \rightarrow \infty$, $n/Y_n \xrightarrow{d} 1$. Hence, as $n \rightarrow \infty$,

$$\begin{aligned} \Lambda(U(X_n))/\Lambda(e^n) &= \exp \left\{ \int_n^{Y_n} G(y) dy \right\} \\ &= \exp \left\{ o(\log[Y_n/n]) \right\} \xrightarrow{d} 1, \end{aligned}$$

and the result follows.

Corollary 2. If $p(s) \equiv 0$ and $\lim_{x \rightarrow \infty} xG(x) = a$, $0 < a < \infty$, then

$$n^{-1} \log U(X_n) - 1 \xrightarrow{d} W,$$

where $P[W \leq u] = \{u/(1+u)\}^a$.

Proof. Immediately, from (6), we have

$$\Lambda(e^{-n}U(X_n))/\Lambda(U(X_n)) \stackrel{a.s.}{\sim} \{1 - n/Y_n\}^a,$$

and the result now follows from Theorem 2.

Corollary 3. If $p(s) \equiv 0$, $\lim_{x \rightarrow \infty} xG(x) = \infty$, and G is regularly varying at infinity, then $nG(\log U(X_n)) \stackrel{d}{\rightarrow} W$, where W is a standard negative exponential random variable.

Proof. Here, from (6), we have

$$\Lambda(e^{-n}U(X_n))/\Lambda(U(X_n)) = \exp \left\{ -Y_n \int_{1-n/Y_n}^1 G(zY_n) dz \right\}; (8)$$

combining (8) with Theorem 2 and $\lim_{x \rightarrow \infty} xG(x) = \infty$, it follows that, as $n \rightarrow \infty$, $n/Y_n \stackrel{d}{\rightarrow} 0$. Hence, since G varies regularly,

$$\Lambda(e^{-n}U(X_n))/\Lambda(U(X_n)) \sim \exp \{-nG(Y_n)\},$$

and the result follows.

If in Corollary 3 G has a positive index Δ , then $0 < \Delta \leq 1$ and the conclusion can be transformed to the form

$$a_n^{-1} \log U(X_n) \stackrel{d}{\rightarrow} W'$$

where W' has the extreme value distribution function $\exp(-x^{-\Delta})$ and $G(a_n) = n^{-1}$.

We now show that for any U the hypotheses of Corollaries 1-3 can be satisfied. Let A be a positive integer valued random variable, define $I = [V(A)]$ and let b be the probability generating function of I . Since

$$V(A) - 1 \leq I \leq V(A) \tag{9}$$

and $\log U$ is SV at infinity, it follows that condition (2) is satis-

fied iff $E \log^+ A < \infty$. In [3, §3.1] several examples were given where $E \log^+ A = \infty$, and for each of these $T(x) = P(A > x)$ is SV at infinity. We now suppose this to be always the case. It follows then, from Lemma 1 and (9), that $P(I > x) \sim T(U(x))$ ($x \rightarrow \infty$) and hence an Abelian theorem for power series yields

$$(1-b(s))/(1-s) = \sum s^j P(I > j) \sim (1-s)^{-1} T(U(1-s)) \quad (s \uparrow 1).$$

Letting $s = \exp(-1/V(e^x))$ and invoking Lemma 1 once again, we obtain

$$G(x) \sim T(e^x). \quad (10)$$

Let $\log_1 x = \log x$ and $\log_k x = \log(\log_{k-1} x)$ ($k=2,3,\dots$) for all sufficiently large x . In [3] an example was given for which

$$T(x) \sim c \left[\prod_{k=1}^r \log_k x \right]^{-1} \quad (x \rightarrow \infty)$$

where $r \geq 2$ and c is a certain constant. Using (10) it is obvious that $\int_0^\infty G(x) dx = \infty$ and that $xG(x) \rightarrow 0$ ($x \rightarrow \infty$) and hence this example satisfies the conditions of Corollary 1.

Discrete distributions were also constructed in [3] for which

$$T(x) \sim a/\log x \quad (0 < a < \infty);$$

$$T(x) \sim c(\log x)^{-(\delta-1)} \quad (0 < c < \infty, 1 < \delta < 2);$$

and

$$T(x) \sim (c/b)(\log_r x)^{-b} \quad (0 < b, c < \infty, r \geq 2).$$

Using (10) we see that these examples satisfy the hypotheses of Corollaries 2 and 3.

Finally, we show that the neatest version of Theorem 2 that could be hoped for is in fact impossible.

Theorem 3. There is no sequence of constants (c_n) such that $c_n^{-1} U(x_n)$ has a limit in distribution which is neither defective nor degenerate at zero.

Proof. For any $0 < x_1 < x_2 < \infty$, let $p_n(x_1, x_2) \equiv P[x_1 \leq c_n^{-1}U(X_n) \leq x_2]$. Then, since $\Lambda(e^{-n}U(\cdot))/\Lambda(U(\cdot))$ is continuous and strictly increasing, it follows also that

$$p_n(x_1, x_2) = P[r_n(x_1) \leq \Lambda(e^{-n}U(X_n))/\Lambda(U(X_n)) \leq r_n(x_2)] ,$$

where

$$0 < r_n(x) \equiv \Lambda(e^{-n}c_n x)/\Lambda(c_n x) < 1.$$

If (c_n) is such that $c_n^{-1}U(X_n)$ is to converge in distribution, we must have $c_n e^{-n} \rightarrow \infty$ because of Theorem 1. Hence, since Λ is SV, we see that $r_n(x_2) = r_n(x_1)[1+o(1)]$ as $n \rightarrow \infty$. Choose any subsequence (n_k) such that $r_{n_k}(x_1) \rightarrow r$ for some $r \in [0, 1]$. Then, for any $\epsilon > 0$, the intervals $[r_{n_k}(x_1), r_{n_k}(x_2)]$ belong to $(r-\epsilon, r+\epsilon)$ for all k sufficiently large. It follows from Theorem 2 that $p_{n_k}(x_1, x_2) \rightarrow 0$, and hence that, if $c_n^{-1}U(X_n)$ converges in distribution, its limit puts no mass on $(0, \infty)$.

It is interesting to note the contrast between the cases $p(x) > 0$ and $p(x) \equiv 0$. In the former, the asymptotic behavior is dominated by the underlying Galton-Watson process, and the effect of immigration, apart from preventing extinction, is seen only in the distribution of the limit of $U(X_n)e^{-n}$: eventually, the contribution of the immigration process becomes negligible. However, when $p(x) \equiv 0$, the immigration distribution has such a broad tail that $U(X_n)e^{-n}$ is pushed off to infinity a.s. by the infinite sequence of occasional, but very large, inflows of immigrants. The character of Theorem 2 and its Corollaries, giving limits in distribution but not with probability one, reflects the nature of the immigration process rather than that of the Galton-Watson process. In particular, unlike the case

when $p(x) > 0$, the limiting distribution appearing in Theorem 2 is the same, whether or not the Galton-Watson process is regular or irregular.

* * * * *

REFERENCES

- [1] Hudson, I. L. and Seneta, E. (1977). The simple branching process with infinite mean. J. Appl. Prob. (to appear)
- [2] Pakes, A. G. (1976). Some limit theorems for a supercritical branching process allowing immigration. J. Appl. Prob., 13, 17-26.
- [3] _____ (1977). Limit theorems for the simple branching process allowing immigration: A review and new results. I: The case of finite offspring mean. Tech. Report No. 128, Series 2, Dept. of Statistics, Princeton University.
- [4] Rényi, A. (1958). On mixing sequences of sets. Acta. Math. Acad. Sci. Hung., 9, 215-228.
- [5] Rényi, A. and Révész, P. (1958). On mixing sequences of random variables. Acta. Math. Acad. Sci. Hung., 9, 389-393.
- [6] Schuh, H-J. and Barbour, A.D. (1977). Normalising constants for the branching process with infinite mean. Adv. Appl. Prob. (to appear)
- [7] Seneta, E. (1976). Regularly Varying Functions. Lecture notes in Mathematics, No. 508. Springer-Verlag, Berlin.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM	
1. REPORT NUMBER Technical Report #134	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER	
4. TITLE (and Subtitle) LIMIT THEOREMS FOR THE SIMPLE BRANCHING PROCESS ALLOWING IMMIGRATION. II. THE CASE OF INFINITE OFFSPRING MEAN.		5. TYPE OF REPORT & PERIOD COVERED Technical rept.	
7. AUTHOR(s) A.D. Barbour, A.D. (Cambridge University) Pakes, A.G.		8. CONTRACT OR GRANT NUMBER(s) 15 N00014-75-C-04534	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Statistics Princeton University, Princeton, N.J.		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS	
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research (Code 436) Arlington, Va. 22217		12. REPORT DATE Dec 1977	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		13. NUMBER OF PAGES 12	
		18. SECURITY CLASS. (of this report) Unclassified	
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.			
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)			
18. SUPPLEMENTARY NOTES Produced while Professor Pakes was serving as a Visiting Research Assistant at Princeton University, on leave from Monash University, Clayton, Victoria, Australia.			
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Bienaymé-Galton-Watson branching process, immigration, martingale convergence, limit theorems, regular variation. sub n (to the -nth power)			
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) This paper obtains some limit theorems for the simple branching process allowing immigration $\langle X_n \rangle$ when the offspring mean is infinite. It is shown that there exists a function U such that $\langle e^{-U(X_n)} \rangle$ converges almost surely and if $s = \sum b_j \log U(j) < \infty$, where $\{b_j\}$ is the immigration distribution, the limit is non-defective and non-degenerate but is infinite if $s = \infty$. When $s = \infty$ limit theorems are found for $\langle U(X_n) \rangle$ which involve a slowly varying non-linear norming. Sum over b sub j log(+) infinity sub j			

DD FORM 1473
1 JAN 73EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-014-8601

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

406 873 JOB